



WHO

Vaccines, Immunization and Biologicals

World Health Organization



English · Español · Français

About us · News · Documents · Statistics, Maps & Charts ·
Site map

Search

VacSurv home ·
V&B home

Last WEB update: September 2004

Vaccine schedule selection form

Schedule for selected antigen: All.

For selected country: Afghanistan

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
* EMR	Afghanistan	BCG	Bacille Calmette-Guérin vaccine	birth;
		DTwP	Diphtheria and tetanus toxoid with whole cell pertussis vaccine	6, 10, 14 weeks;
		Measles	Measles vaccine	9 months;
		OPV	Oral polio vaccine	birth; 14 days; 6, 10, 14 weeks; 9 months;
		TT	Tetanus toxoid	15-45 years; +1, +6 months; +1, +1 year;

To contact us:

World Health Organization

Department of Immunization Vaccines and Biologicals, Vaccine Assessment and Monitoring Team
CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

Archives · Contact WHO · © 2004 WHO/OMS · Search WHO Web site · Search other UN Web sites · Suggest



WHO.int

Vaccines, Immunization and Biologicals

World Health Organization



English · Español · Français

About us · News · Documents · Statistics, Maps & Charts ·
Site map

Search

VacSurv home ·
V&B home

Last WEB update: September 2004 Vaccine schedule selection form

Schedule for selected antigen: All.

For selected country: Argentina

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
AMR	Argentina	BCG	Bacille Calmette-Guérin vaccine	birth; 6 years;
		DTP	Diphtheria and tetanus toxoid with pertussis vaccine	6 years;
		DTPHiB	Tetavalent diphtheria and tetanus toxoid with pertussis and Hib vaccine	2, 4, 6, 18 months;
		HepB	Hepatitis B vaccine	birth; 2, 6 months;
		MMR	Measles mumps and rubella vaccine	12 months;
		OPV	Oral polio vaccine	2, 4, 6, 18 months; 6 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	>16 years; and pregnant women

To contact us:

World Health Organization

Department of Immunization Vaccines and Biologicals, Vaccine Assessment and Monitoring Team

CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

Archives · Contact WHO · © 2004 WHO/OMS · Search WHO Web site · Search other UN Web sites · Sugg



WHO:INT

World Health Organization

Vaccines, Immunization and Biologicals


[English](#) · [Español](#) · [Français](#)
[About us](#) · [News](#) · [Documents](#) · [Statistics, Maps & Charts](#) · [Site map](#)
[Search](#)
[VacSurv home](#) · [V&B home](#)
Schedule for selected antigen: All.
For selected country: Bosnia and Herzegovina

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
EUR	Bosnia & Herzegovina	BCG	Bacille Calmette-Guérin vaccine	birth;
		DT	Tetanus and diphtheria toxoid childrens' dose	7 years;
		DTWP	Diphtheria and tetanus toxoid with whole cell pertussis vaccine	2, 4, 6 months; 2, 4 years;
		HepB	Hepatitis B vaccine	7 years;
		MMR	Measles mumps and rubella vaccine	2, 7 years;
		OPV	Oral polio vaccine	2, 4, 6 months; 2, 4 years; (7 years, 14 years)
		Rubella	Rubella vaccine	14 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	14 years;
		TT	Tetanus toxoid	18 years;

To contact us:

World Health Organization

Department of Vaccines and Biologicals Vaccine Assessment and Monitoring Team

CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227



WHO.INT

World Health Organization

Vaccines, Immunization and Biologicals



English · Español · Français

Search

[About us](#) · [News](#) · [Documents](#) · [Statistics, Maps & Charts](#) · [Site map](#)

[VacSurv home](#) · [V&B home](#)

Schedule for selected antigen: All.
For selected country: Costa Rica

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
AMR	Costa Rica	BCG	Bacille Calmette-Guérin vaccine	birth;
		DT	Tetanus and diphtheria toxoid childrens' dose	school; adult x2
		DTP	Diphtheria and tetanus toxoid with pertussis vaccine	2, 4, 6, 15 months;
		HepB	Hepatitis B vaccine	birth, 2, 6 months;
		Hib	Haemophilus influenzae type b vaccine	2, 4, 6, 15 months;
		MMR	Measles mumps and rubella vaccine	15 months; 6 years;
		OPV	Oral polio vaccine	2, 4, 6 months; 4 years;

To contact us:
 World Health Organization
 Department of Vaccines and Biologicals Vaccine Assessment and Monitoring Team
 CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227



WHO.int

World Health Organization

Vaccines, Immunization and Biologicals



English · Español · Français

About us · News · Documents · Statistics, Maps & Charts · Site map

Search
 VacSurv home · V&B home

Schedule for selected antigen: All.

For selected country: Ethiopia

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
AFR	Ethiopia	BCG	Bacille Calmette-Guérin vaccine	birth;
		DTwP	Diphtheria and tetanus toxoid with whole cell pertussis vaccine	6, 10, 14 weeks;
		Measles	Measles vaccine	+9 months;
		OPV	Oral polio vaccine	6, 10, 14 weeks;
		TT	Tetanus toxoid	first contact; +1, +6 months; +1 year; CBAW 15-49 years
		VitaminA	Vitamin A supplementation	9 months;

To contact us:
 World Health Organization
 Department of Vaccines and Biologicals Vaccine Assessment and Monitoring Team
 CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

Archives · Contact WHO · © 2004 WHO/OMS · Search WHO Web site · Search other UN Web sites · Suggest



WHO.INT

World Health Organization

Vaccines, Immunization and Biologicals



English · Español · Français

[About us](#) · [News](#) · [Documents](#) · [Statistics, Maps & Charts](#) · [Site map](#)

Search

[VacSurv home](#) · [V&B home](#)

Schedule for selected antigen: All.

For selected country: Iraq

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
EMR	Iraq	BCG	Bacille Calmette-Guérin vaccine	birth
		DT	Tetanus and diphtheria toxoid childrens' dose	<5 years;
		DTP	Diphtheria and tetanus toxoid with pertussis vaccine	2, 4, 6, 18 months, 4- 6 years;
		HepB	Hepatitis B vaccine	birth, 2, 6 months;
		Measles	Measles vaccine	9 months;
		MMR	Measles mumps and rubella vaccine	school entry;
		OPV	Oral polio vaccine	birth, 2, 4, 6, 18 months, 4-6 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	7 years;
		TT	Tetanus toxoid	CBAW x5;

To contact us:

World Health Organization

Department of Vaccines and Biologicals Vaccine Assessment and Monitoring Team

CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227



WHO/INT

Vaccines, Immunization
and Biologicals

World Health Organization



English · Español · Français

About us · News · Documents · Statistics, Maps & Charts ·
Site map.

Search

VacSurv home ·
V&B home

Last WEB update: September 2004

Vaccine schedule selection form

Schedule for selected antigen: All.

For selected country: Iran (Islamic Republic of)

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
EMR	Iran	BCG	Bacille Calmette-Guérin vaccine	birth;
		DTwP	Diphtheria and tetanus toxoid with whole cell pertussis vaccine	6, 12, 18 weeks; 15 months; 4-6 years;
		HepB	Hepatitis B vaccine	birth; 6 weeks; 9 months;
		Measles	Measles vaccine	9, 15 months;
		OPV	Oral polio vaccine	birth; 6, 12, 18 weeks; 15 months; 4-6 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	14-16 years;

To contact us:

World Health Organization

Department of Immunization Vaccines and Biologicals, Vaccine Assessment and Monitoring Team

CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

Archives · Contact WHO · © 2004 WHO/OMS · Search WHO Web site · Search other UN Web sites · Sugg



WHO

Vaccines, Immunization and Biologicals

World Health Organization



English · Español · Français

About us · News · Documents · Statistics, Maps & Charts ·
Site map

Search

VacSurv home ·
V&B home

Last WEB update: September 2004 Vaccine schedule selection form

Schedule for selected antigen: All.

For selected country: Israel

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
EUR	Israel	DTaP	Diphtheria and tetanus toxoid with acellular pertussis vaccine	2, 4, 6 months; 1 year;
		HepA	Hepatitis A vaccine	18, 24 months;
		HepB	Hepatitis B vaccine	birth; 1, 6 months;
		Hib	Haemophilus influenzae type b vaccine	2, 4, 6 months; 1 year;
		IPV	Inactivated polio vaccine	2, 4, 12 months;
		MMR	Measles mumps and rubella vaccine	12 months; 6 years;
		OPV	Oral polio vaccine	4, 6, 12 months; 6, 13 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	8-9, 13-14 years;

To contact us:

World Health Organization

Department of Immunization Vaccines and Biologicals, Vaccine Assessment and Monitoring Team

CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

Archives · Contact WHO · © 2004 WHO/OMS · Search WHO Web site · Search other UN Web sites · Sugg



Vaccines, Immunization and Biologicals



English · Español · Français

About us · News · Documents · Statistics, Maps & Charts · Site map

Search

VacSurv home · V&B home

Schedule for selected antigen: All.

For selected country: Jordan

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
EMR	Jordan	BCG	Bacille Calmette-Guérin vaccine	birth; (vaccination started in some health centers)
		DT	Tetanus and diphtheria toxoid childrens' dose	2, 3, 4, 18 months; (some cases)
		DTPHep	Tetavalent diphtheria and tetanus toxoid with pertussis and HepB vaccine	2, 3, 4 months; (in the first half of the year)
		DTPHib	Tetavalent diphtheria and tetanus toxoid with pertussis and Hib vaccine	2, 3, 4 months; (in the first half of the year)
		DTwP	Diphtheria and tetanus toxoid with whole cell pertussis vaccine	18 months;
		HepB	Hepatitis B vaccine	2, 3, 4 months; (in the first half of the year)
		Hib	Haemophilus influenzae type b vaccine	2, 3, 4 months; (in the first half of the year)
		Measles	Measles vaccine	9 months;
		MMR	Measles mumps and rubella vaccine	18 months;
		OPV	Oral polio vaccine	2; 3, 4, 9, 18 months; 6 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	6, 15 years; (school children)
		TT	Tetanus toxoid	first contact; +1, +6 months; +1, +1 year;

To contact us:

World Health Organization

Department of Vaccines and Biologicals Vaccine Assessment and Monitoring Team

CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227



WHO.int

World Health Organization

Vaccines, Immunization and Biologicals



English · Español · Français

About us · News · Documents · Statistics, Maps & Charts · Site map

Search

VacSurv home · V&B home

Last WEB update: September 2004 Vaccine schedule selection form

Schedule for selected antigen: All.

For selected country: Lebanon

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
EMR	Lebanon	DTwP	Diphtheria and tetanus toxoid with whole cell pertussis vaccine	4-5 years;
		DTwPHib	Diphtheria and tetanus toxoid with whole cell pertussis and Hib vaccine	2, 3, 4, 15-18 months;
		HepB	Hepatitis B vaccine	birth; 1, 4 months;
		MMR	Measles mumps and rubella vaccine	13 months; 4-5 years;
		OPV	Oral polio vaccine	2, 3, 4, 15-18 months; 4-5, 10-12 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	10-12 years;

To contact us:

World Health Organization

Department of Immunization Vaccines and Biologicals, Vaccine Assessment and Monitoring Team
 CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

Archives · Contact WHO · © 2004 WHO/OMS · Search WHO Web site · Search other UN Web sites · Suggest



VACCINES

World Health Organization

Vaccines, Immunization
and Biologicals

English · Español · Français

[About us](#) · [News](#) · [Documents](#) · [Statistics, Maps & Charts](#) · [Site map](#)[Search](#)[VacSurv home](#) · [V&B home](#)

Last WEB update: September, 2004

Vaccine schedule selection form

Schedule for selected antigen: All.

For selected country: Poland

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
EUR	Poland	BCG	Bacille Calmette-Guérin vaccine	birth; 12* months; 7, 12 years; *only if no scar developed
		DT	Tetanus and diphtheria toxoid childrens' dose	6 years; until Sept. 2003
		DTaP	Diphtheria and tetanus toxoid with acellular pertussis vaccine	6 years; since Oct. 2003
		DTwP	Diphtheria and tetanus toxoid with whole cell pertussis vaccine	2, 3-4, 5, 16-18 months;
		HepB	Hepatitis B vaccine	birth; 2, 6-7 months;
		IPV	Inactivated polio vaccine	3-4, 5 months;
		Measles	Measles vaccine	13-14 months; 7 years; until Oct. 2003
		MMR	Measles mumps and rubella vaccine	13-14 months; since Nov. 2003
		OPV	Oral polio vaccine	6-7, 16-18 months; 6 years;
		Rubella	Rubella vaccine	13 years; female
		Td	Tetanus and diphtheria toxoid for older children / adults	14, 19 years;

To contact us:

World Health Organization

Department of Immunization Vaccines and Biologicals, Vaccine Assessment and Monitoring Team

CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

[Archives](#) · [Contact WHO](#) · [© 2004 WHO/OMS](#) · [Search WHO Web site](#) · [Search other UN Web sites](#) · [Sugg](#)



WORLD HEALTH ORGANIZATION

World Health Organization

Vaccines, Immunization and Biologicals



English · Español · Français

About us · News · Documents · Statistics, Maps & Charts · Site map

Search
VacSurv home · V&B home

Last WEB update: September 2004 Vaccine schedule selection form

Schedule for selected antigen: All.

For selected country: Serbia and Montenegro

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
EUR	Serbia & Montenegro	BCG	Bacille Calmette-Guérin vaccine	birth;
		DT	Tetanus and diphtheria toxoid childrens' dose	7 years;
		DTWP	Diphtheria and tetanus toxoid with whole cell pertussis vaccine	8, 14, 20 weeks;
		HepB	Hepatitis B vaccine	8, 14 weeks; 12 months;
		MMR	Measles mumps and rubella vaccine	12 months; 12 years;
		OPV	Oral polio vaccine	8, 14, 20 weeks; 15 months; 7, 14 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	14 years;
		TT	Tetanus toxoid	18 years;

To contact us:

World Health Organization
Department of Immunization Vaccines and Biologicals, Vaccine Assessment and Monitoring Team
CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

Archives · Contact WHO · © 2004 WHO/OMS · Search WHO Web site · Search other UN Web sites · Suggest



WHO/INT

World Health Organization

Vaccines, Immunization and Biologicals



English · Español · Français

About us · News · Documents · Statistics, Maps & Charts · Site map

Search

VacSurv home · V&B home

Last WEB update: September 2004 Vaccine schedule selection form

Schedule for selected antigen: All.

For selected country: Syrian Arab Republic

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
EMR	Syria	BCG	Bacille Calmette-Guérin vaccine	birth;
		DT	Tetanus and diphtheria toxoid childrens' dose	6 years;
		DTwPHib	Diphtheria and tetanus toxoid with whole cell pertussis and Hib vaccine	3, 4, 5 months;
		HepB	Hepatitis B vaccine	birth; 3, 10 months;
		Measles	Measles vaccine	10 months;
		MMR	Measles mumps and rubella vaccine	15 months;
		OPV	Oral polio vaccine	birth; 3, 4, 5, 18 months; 6 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	12 years;
		TT	Tetanus toxoid	CBAW;
		VitaminA	Vitamin A supplementation	10 months;

To contact us:

World Health Organization

Department of Immunization Vaccines and Biologicals, Vaccine Assessment and Monitoring Team

CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

Archives · Contact WHO · © 2004 WHO/OMS · Search WHO Web site · Search other UN Web sites · Suggest



WHO

World Health Organization

Vaccines, Immunization and Biologicals



English · Español · Français

About us · News · Documents · Statistics, Maps & Charts ·
Site map

Search
VacSurv home ·
V&B home

Schedule for selected antigen: All.

For selected country: Thailand

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
SEAR	Thailand	BCG	Bacille Calmette-Guérin vaccine	birth;
		DTPHep	Tetavalent diphtheria and tetanus toxoid with pertussis and HepB vaccine	2, 4, 6 months; (only in Chiang Rai Province)
		DTwP	Diphtheria and tetanus toxoid with whole cell pertussis vaccine	2, 4, 6 months; 1.5-2, 4-5 years;
		HepB	Hepatitis B vaccine	birth; 2, 6 months;
		JapEnc	Japanese encephalitis	1.5-2, 1.5-2, 2.5-3 years;
		Measles	Measles vaccine	9-12 months;
		MMR	Measles mumps and rubella vaccine	6 years; (school children 1st grade)
		OPV	Oral polio vaccine	2, 4, 6 months; 1.5-2, 4-5 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	6, 12 years;
		TT	Tetanus toxoid	pregnant women; first contact; +1, +6 months; CBAW 15-45 years (only in high risk area)

To contact us:

World Health Organization

Department of Vaccines and Biologicals Vaccine Assessment and Monitoring Team

CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

Archives · Contact WHO · © 2004 WHO/OMS · Search WHO Web site · Search other UN Web sites · Sugg



WHO/INT

Vaccines, Immunization and Biologicals

World Health Organization



[English](#) · [Español](#) · [Français](#)

[About us](#) · [News](#) · [Documents](#) · [Statistics, Maps & Charts](#) · [Site map](#)

[Search](#)

[VacSurv home](#) · [V&B home](#)

Schedule for selected antigen: All.

For selected country: Turkey

Region	Country	Antigen (or nutritional supplement)	Description	Schedule
EUR	Turkey	BCG	Bacille Calmette-Guérin vaccine	2 months; 6 years;
		DTwP	Diphtheria and tetanus toxoid with whole cell pertussis vaccine	2, 3, 4, 16-24 months;
		HepB	Hepatitis B vaccine	3, 3, 4, 9 months;
		Measles	Measles vaccine	9 months; 6 years;
		OPV	Oral polio vaccine	2, 3, 4 months; 6 years;
		Td	Tetanus and diphtheria toxoid for older children / adults	6, 11 year;
		TT	Tetanus toxoid	180, 181, 187, 199, 211 months;

To contact us:

World Health Organization

Department of Vaccines and Biologicals Vaccine Assessment and Monitoring Team

CH-1211 Geneva 27, Switzerland. Fax: 41 22 791 4227

[Archives](#) · [Contact WHO](#) · [© 2004 WHO/OMS](#) · [Search WHO Web site](#) · [Search other UN Web sites](#) · [Sugg](#)